

FIPROTEC[®]



spray



INDICATIONS:

- For the treatment and prevention of flea and tick infestation on dogs and aids in the control of flea allergic dermatitis on dogs.
- Long acting flea and tick treatment for dogs.
- FIPROTEC[®] spray may be used on pregnant and lactating females and on unweaned puppies.

COMPOSITION:

Fipronil 0.25 % m/v (0.25g/100ml)

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- DO NOT bath or shampoo the animal 2 days before or 2 days after the application as this will decrease the duration of activity.
- Ruffle the dry coat against the lie of the hair before spraying.
- Spray the entire body of the animal at a distance of 10 - 20 cm.
- To ensure coverage of the head region spray FIPROTEC[®] spray onto a glove and rub gently around the face, avoiding the eyes.
- When your pet has been thoroughly covered, massage the coat all over to ensure that the spray gets right down to the skin.
- When treating puppies you may prefer to use a glove to apply FIPROTEC[®] spray.
- Allow your pet to dry naturally. Do not towel dry.
- Monthly applications are recommended where there is a high risk of re-infestation, if the pet is allergic to fleas and in situations where tick control is needed.
- For this control to be effective all dogs and cats in the household should be treated simultaneously.

APPLICATION:

In order to wet the coat down to the skin it is recommended that the following application rates be used:

Short-haired animals (<1.5 cm)

- A minimum of **3 ml/kg** body mass = 7,5 mg of active material kg/body mass.

Long-haired animals (>1.5 cm)

- A minimum of **6 ml/kg** body mass = 15 mg of active material kg/body mass.

100 ml bottle

Each trigger application delivers 0.5 ml spray volume

e.g. for dogs under 12 kg

6 pumps actions per kg

Weight 5 kg = 30 pump actions

Weight 10 kg = 60 pump actions

250 ml bottle

Each trigger application delivers 1 ml spray volume

e.g. for large dogs over 12 kg

3 pump actions per kg

Weight 15 kg = 45 pumps actions

Weight 30 kg = 90 pumps actions

RESIDUAL EFFECT

• Fleas: 2 months (1 - 3 months)

• Ticks: 1 month (3 - 5 weeks)

This will depend on the parasite challenge in the animal's environment

WARNINGS:

- Keep out of reach of uninformed persons, children and animals
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS:

- Toxic to bees and fish
- Avoid contact with eyes
- Wear gloves, especially during treatment of groups of animals
- Spray in a well ventilated area
- Wash hands immediately after use.
- Do not smoke, eat or drink during application.
- Keep your pet away from fires, heat or surfaces likely to be affected by the alcohol spray until dry.

PRESENTATION

100ml and 250ml white plastic spray bottle

STORAGE INSTRUCTIONS:

Store in a cool place.

REGISTRATION NUMBER: G3793 (Act 36/1947)

REGISTRATION HOLDER:

Cipla Vet (Pty) Ltd.
Co. Reg. No. 2001/017471/07
PO Box 1096, Durbanville, 7551
Tel: 0861 115 037
Fax: 0861 115 038

FOR FULL PARTICULARS SEE ENCLOSED PACKAGE INSERT.